arXiv:1612.09267v1 [cond-mat.mes-hall] 29 Dec 2016

Amorphous gyroscopic topological metamaterials

Noah P. Mitchell,[1,*] Lisa M. Nash,[1] Daniel Hexner,[1] Ari Turner,[2] and William T. M. Irvine[1,3,†]

[1] *James Franck Institute and Department of Physics, University of Chicago, Chicago, IL 60637, USA*
[2] *Department of Physics, Israel Institute of Technology*
[3] *Enrico Fermi Institute, The University of Chicago, Chicago, IL 60637, USA*

The discovery that the band structure of electronic insulators may be topologically non-trivial has unveiled distinct phases of electronic matter with novel properties [4, 7]. Recently, mechanical lattices have been found to have similarly rich structure in their phononic excitations [6, 15], giving rise to protected uni-directional edge modes whose existence was demonstrated in lattices of interacting gyroscopes and coupled pendula [11, 19, 23]. In all these cases, however, as well as in other topological metamaterials [6, 18], the underlying structure was finely tuned, be it through periodicity, quasi-periodicity or isostaticity. Here we show that amorphous mechanical Chern insulators consisting of interacting gyroscopes can be readily constructed from arbitrary underlying structures, including hyperuniform, jammed, quasi-crystalline, and uniformly random arrangements. We find that local geometry and local interactions control the topology of the vibrational spectrum, allowing simple, local decorations to endow amorphous structures with protected edge modes—with a chirality of choice. Using a real-space generalization of the Chern number, we investigate the topology of our structures numerically, analytically and experimentally. The robustness of our approach enables the topological design and self-assembly of non-crystalline topological metamaterials on the micro and macro scale.

The quintessential model for the mechanical behavior of materials is a network of masses connected by springs. Though such a material is simple in its constituents, varying its connectivity can generate a plethora of interesting and desirable mechanical properties. While the search for new materials within this paradigm continues [6, 13, 18], the possibility of replacing the masses with active constituent particles allows new material behaviors not possible in the mass-and-spring paradigm [2, 10, 17].

A recent example found that introducing active chiral components using rapidly spinning tops (or 'gyroscopes') gives rise to a simple model system that breaks time-reversal symmetry [11, 23]. The system consists of gyroscopes suspended from a plate that interact via springs or magnetically. The response of a gyroscope to an applied force is fundamentally different than that of a mass: it is first order in time and perpendicular to the applied force. Because of this peculiar response, lattices of gyroscopes with a spring-like interaction obey a Schrödinger-like equation. Appropriate crystalline arrangements of these gyroscopes were found to break time-reversal symmetry, opening topological phononic band gaps and supporting robust chiral edge modes [11].

Unlike trivial insulators, whose electronic states can be thought of as a sum of independent local insulating states, topological insulators have necessarily longer-range correlations [20, 22]. It is natural, therefore, to assume that some regularity over long distances is key to topological behavior, even if topological properties are robust to the addition of disorder. However, the extent to which spatial order needs to be built into the structure that gives rise to topological modes is unclear. We report a recipe for constructing amorphous arrangements of interacting gyroscopes—structurally more akin to a liquid than a solid—that naturally support topological phonon spectra. This shows that topology can naturally arise in materials for which the only design principle is the local connectivity, a criterion that lends it self to imperfect manufacturing and self-assembly.

Starting from an arbitrary point set such as the one shown in Figure 1a, a natural way to form a network is to generate a Voronoi tessellation, either via the Wigner-Seitz construction or by connecting centroids of a triangulation [3]. Treating the edges of the cells as bonds and placing gyroscopes at the vertices leads to a network reminiscent of 'topological disorder' in electronic systems [21]. The top panel in Figure 3a shows the resulting overlaid density of states for ten 'Voronoized' hyperuniform networks, colored by the localization length of the mode, λ. A range of frequencies arises in which all modes are tightly localized, and this frequency region overlaps with the corresponding band gap of the honeycomb lattice. We find that gyroscope-and-spring networks constructed in this way from arbitrary initial point sets invariably have such a mobility gap in an frequency range determined by the strength of the gravitational pinning and spring interactions.

Our networks are structurally akin to liquids and thus the essential features of the energy spectrum described above emerge in the complete absence of long range structural order. Furthermore, the details of the Voronoization are not essential either, as can be verified by replacing the centroidal construction [3] with a Wigner-Seitz construction (see Supplementary Information Figures 21-23 for a comparison).

Our networks are reminiscent of 'topologically disordered' electronic systems [9]. In these systems, a central characteristic is that the local density of states as a function of frequency is predictive of the global density of states. Specifically, band gaps or mobility gaps are pre-



FIG. 1. **Decorations of an amorphous structure preserve isotropy and lack of long-range order** | Two-point correlation functions $g(\mathbf{x})$ (right) are shown for a system of $N \approx 3000$ to 6000 particles; the number of particles increases with each decoration. **a**, A snippet of a hyperuniform point set. **b**, Voronoi decoration of the point set, constructed by connecting adjacent centroids of a triangulation. **c**, Placing gyroscopes at the midpoints of the Voronoi network's bonds results in the Kagome decoration.

served [5, 24, 25]. Interestingly, we find that, even in the presence of band topology, averaging the local density of states over mesoscopic patches (~ 10 gyroscopes) reproduces the essential features of the global density of states as a function of frequency. Furthermore, we find that inserting mesoscopic patches of our structures into a variety of other dissimilar networks (see Supplementary Information Figures 8-10) does not significantly disrupt the averaged local density of states of the patch.

Crucially, we find that our structures show hallmarks of non-trivial topology. When the system is cut to a finite size, modes confined to the edge populate the mobility gap, mixed in with localized states. As shown in the direct simulations of Figure 2 and Supplementary Videos 1 and 2, shaking a gyroscope on the boundary results in chiral waves that bear all the hallmarks of protected edge states (robustness to disorder and absence of backscattering).

An experimental realization can be readily constructed from gyroscopes interacting magnetically, as shown in Figure 2. Like in [11], these gyroscopes are constructed of 3D-printed units encasing DC motors which interact via magnetic repulsion. Probing the edge of this system immediately generates a chiral wave packet localized to the boundary, confirming that this class of topological material is physically realizable and robust (see Figure 2c and Supplementary Video 3).

This behavior begs for a topological characterization, even though it might be surprising that topology can emerge from such a local construction. The existence of chiral edge states in an energy gap is guaranteed if an invariant known as the Chern number is nonzero, and the direction of the chiral waves is given by its sign. Although the Chern number was originally defined in momentum space, several generalizations have been constructed in coordinate space in order accommodate disorder in crystalline and quasi-crystalline electronic materials [1, 8, 14]. In these methods, information about the system's vibrations is carried by the projection operator, P. Each element P_{ij} measures the response of gyroscope j to excitations of gyroscope i within a prescribed range (band) of frequencies.

According to one such method, proposed in [8], a subset of the system is divided into three parts and labeled in a counterclockwise fashion (red, green, and blue regions in Figure 3). These regions are then used to index components of an antisymmetric product of projection operators:

$$\nu(P) = 12\pi i \sum_{j\in A}\sum_{k\in B}\sum_{l\in C}\left(P_{jk}P_{kl}P_{lj} - P_{jl}P_{lk}P_{kj}\right). \quad (1)$$

The sum of such elements converges to the Chern number of the chosen band after the region has enclosed a few lattice spacings (see Supplementary Videos 4 and 5 and Supplementary Information).

Equation 1 can be understood as a form of charge polarization in the response of an electronic material to a locally applied magnetic field. Applying a magnetic field to a small region of a material induces an electromotive force winding around the site of application. If the material is a trivial insulator, any changes in charge density there arise from local charge re-arrangements, which result in no accumulation of charge. By contrast, a topological electronic system has a Hall conductivity determined by the Chern number. As a result, a net current will flow perpendicular to the electromotive force, inducing a net-nonzero charge concentrated at the magnetic field site, compensated by charge on the boundary. As we show



FIG. 2. **Time-domain simulations and experiments confirm chiral edge modes** | Voronoization (**a**) and kagomization (**b**) of an amorphous point set results in topological gyroscopic networks with chiral edge modes propagating in opposing directions. The networks shown here are based on hyperuniform point sets, and localization of modes is probed by participation ratio, $p = \left(\sum_i |\psi_i|^2\right)^2 / N \sum_i |\psi_i|^4$, since the edge modes are not exponentially localized. **c**, Edge mode propagating in a amorphous experimental gyroscopic network based on a portion of a jammed network configuration. The motor-driven gyroscopes couple via a magnetic dipole-dipole interaction. Despite the nonlinear interaction and spinning speed disorder ($\sim 10\%$), the edge mode persists.

in the Supplementary Information, the amount of local charging is proportional to the applied field, and the proportionality constant is the Chern number of Equation 1.

Figure 3a shows the results computed for the Voronoized networks. As the cutoff frequency for the projector is varied, the Chern number converges to $\nu = -1$ in a broad range of frequencies centered about the mobility gap, confirming that the modes observed in Figure 2 are topological in origin and predicting their direction.

Having established this connection, it is natural to ask whether the sign of the Chern number can be controlled. What in our network construction gives rise to this value? What amorphous networks have an upper band with a Chern number of $\nu = 0$ or $+1$? To investigate this point, we construct alternative decorations of the same initial point set. One possible construction arises naturally from



FIG. 3. **Chern spectra confirm topological mobility gaps** | The Chern number is computed for the band of frequencies above ω_c using a real-space method. Overlaid density of states $D(\omega)$ histograms for ten realizations of (**a**) Voronoized hyperuniform point sets ($\sim$ 2000 particles) and (**b**) kagomized hyperuniform point sets ($\sim$ 2500 particles), with each mode colored by its localization length, λ. The topological mobility gaps remain in place and populated by a relatively small number of highly localized states for all realizations. All networks have their precession and spring frequencies set to be equal ($\Omega_g = \Omega_k$).

joining neighboring points in the original point set, leading to a Delaunay triangulation. As shown in Supplementary Video 6, such networks show no gaps and no topology, suggesting that the local geometry dictated by Voronoization is responsible for the emergent topology. A clue can be found by noting that the Voronoized networks are locally akin to a honeycomb lattice. This is the simplest lattice with two sites per unit cell, a necessary condition for supporting a band gap. Moreover, this lattice was previously found to be topological with the same Chern number [11].

Building on this insight, we introduce a second decoration, which we dub 'kagomization,' shown in Figure 1c. If applied to a triangular point set, Voronoization produces a honeycomb lattice and kagomization produces a kagome lattice, the simplest lattice with three sites per unit cell which we have found to produce $\nu = +1$ gyroscopic metamaterials. Proceeding as with the Voronoized network case (Figures 2 and 3), we find the presence of topologically protected modes with opposite direction and the corresponding opposite Chern number in the the band structure (see Supplementary Video 7).

The dependence of the Chern number on *local* decorations of random point sets suggests that the topology of the vibrational spectrum in these structures is local in origin. A strong confirmation of this insight comes from considering the projection operator P. The projector value P_{ij} measures the vibrational correlation between gyroscope j and gyroscope i when considering all modes above a cutoff frequency. By explicitly computing its magnitude in our networks, we find that the magnitude of P_{ij} falls off exponentially with distance (Supplementary Information Figure 12). Remarkably, explicitly cutting out a section of the network and embedding it in a network with a different spectrum results in only a slight change to the local projector values ($< 2\%$) (see Supplementary Information Figure 13). Since the Chern number is built from these nearest-neighbor projector elements, it then follows that the local structure of the gyroscope network, combined with some homogeneity of this local structure across the lattice, is all that is needed to determine the Chern number (c.f. Supplementary Video 8).

This situation is reminiscent of electronic glasses in which the local binding structure gives rise to a local 'gap.' Under weak assumptions of homogeneity, this gap can be shown to extend to the whole system [24, 25]. The case with topology is similar: the next-nearest neighbor angles in a network's cell open a local 'gap' via time reversal symmetry breaking.

As in the lattice case, a mobility gap becomes topological due to time reversal symmetry breaking: bond angles in these networks are not multiples of $\pi/2$. We can probe this mechanism by eliminating a gap's topology. Alternating the pinning strength, Ω_g, of neighboring gyroscopes in a network mimics the breaking of inversion symmetry on a local scale, an effect which competes against the time reversal gap opening mechanism. When the pinning strength difference between sites is large enough, this competing mechanism eliminates edge modes, triggering a transition to a $\nu = 0$ mobility gap, shown in Figure 4a and Supplementary Videos 9 and 10.



FIG. 4. **Transition of a topological amorphous network to the trivial phase, and binary mixtures of Voronoized and kagomized networks** | **a**, Locally breaking inversion symmetry by increasing and decreasing the precession frequencies of alternating gyroscopes competes with broken time reversal symmetry, triggering a transition to the trivial phase, with no edge modes. The precession frequency splitting, Δ, is tuned so that $\Omega_g^A = \Omega_k(1+\Delta)$ and $\Omega_g^B = \Omega_k(1-\Delta)$. **b**, Edge modes are localized at the interfaces between kagomized and Voronoized networks, permitting sinuous channels for the propagation of unidirectional phonons. **c**, Excitations of a Voronoized region nested inside a kagomized network remain confined when the excitation frequency is in a mobility gap unique to the kagomized network. **d**, When kagomized elements are randomly mixed into a Voronoized network, the sign of the local, spatially-resolved Chern calculation is determined by the local geometry, with excitations in a mobility gap biased toward the interface of the two clusters.

Equipped with these insights, we can easily engineer networks which are heterogeneous mixtures of these two local configurations. Figure 4b-d highlight some results of combining both decorations or encapsulating one within another. Because the Voronoized and kagomized networks share a mobility gap, excitations are localized to their interface, offering a method of creating robust unidirectional waveguides, such as the sinuous waveguide shown in Figure 4b and Supplementary Video 11. Figure 4c demonstrates that additional topological mobility gaps at higher frequency in the kagomized network allow bulk excitations to be confined to an encapsulated Voronoized region (see also Supplementary Video 12). Random mixtures of the two decorations, shown in Figure4d, demonstrate heterogeneous local Chern number measurements, with mobility-gap excitations biased toward the interfaces between red and blue regions (see Supplementary Information Figure 14 and Supplemen-

tary Video 13).

Our study demonstrates that local interactions and local geometric arrangement are sufficient to generate chiral edge modes on the scale of the system size, promising new avenues the engineering of topological mechanical metamaterials generated via imperfect self-assembly processes. Such self-assembled materials could be constructed, for instance, with micron-scale spinning magnetic particles. Since our methods bear substantial resemblance to electronic tight-binding models, our results also find application in the study of topological electronic materials, photonic Floquet topological insulators [16], and coupled RC circuits [12].

* npmitchell@uchicago.edu; Corresponding author
† wtmirvine@uchicago.edu; Corresponding author

[1] Raffaello Bianco and Raffaele Resta. Mapping topological order in coordinate space. *Physical Review B*, 84(24):241106, December 2011.

[2] Romain Fleury, Alexander B Khanikaev, and Andrea Alú. Floquet topological insulators for sound. *Nature Communications*, 7:11744, June 2016.

[3] Marian Florescu, Salvatore Torquato, and Paul J. Steinhardt. Designer disordered materials with large, complete photonic band gaps. *Proceedings of the National Academy of Sciences*, 106(49):20658–20663, December 2009.

[4] F.D.M. Haldane. Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the "Parity Anomaly". *Phys Rev Lett*, 61(18):2015–2018, October 1988.

[5] R. Haydock, V. Heine, and M. J. Kelly. Electronic structure based on the local atomic environment for tight-binding bands. *Journal of Physics C: Solid State Physics*, 5(20):2845, 1972.

[6] C. L. Kane and T. C. Lubensky. Topological boundary modes in isostatic lattices. *Nat Phys*, 10(1):39–45, December 2013.

[7] C. L. Kane and E. J. Mele. Quantum Spin Hall Effect in Graphene. *Physical Review Letters*, 95(22):226801, November 2005.

[8] Alexei Kitaev. Anyons in an exactly solved model and beyond. *Annals of Physics*, 321(1):2–111, January 2006.

[9] B. Kramer and A. MacKinnon. Localization: theory and experiment. *Reports on Progress in Physics*, 56(12):1469, 1993.

[10] M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, Madan Rao, and R. Aditi Simha. Hydrodynamics of soft active matter. *Reviews of Modern Physics*, 85(3):1143–1189, July 2013.

[11] Lisa M. Nash, Dustin Kleckner, Alismari Read, Vincenzo Vitelli, Ari M. Turner, and William T. M. Irvine. Topological mechanics of gyroscopic metamaterials. *Proceedings of the National Academy of Sciences*, 112(47):14495–14500, November 2015.

[12] Jia Ningyuan, Clai Owens, Ariel Sommer, David Schuster, and Jonathan Simon. Time- and Site-Resolved Dynamics in a Topological Circuit. *Phys Rev X*, 5(2):021031, June 2015.

[13] Martin Ostoja-Starzewski. Lattice models in micromechanics. *Applied Mechanics Reviews*, 55(1):35, 2002.

[14] Emil Prodan. Non-commutative tools for topological insulators. *New Journal of Physics*, 12(6):065003, 2010.

[15] Emil Prodan and Camelia Prodan. Topological Phonon Modes and Their Role in Dynamic Instability of Microtubules. *Phys Rev Lett*, 103(24):248101, December 2009.

[16] Mikael C. Rechtsman, Julia M. Zeuner, Yonatan Plotnik, Yaakov Lumer, Daniel Podolsky, Felix Dreisow, Stefan Nolte, Mordechai Segev, and Alexander Szameit. Photonic Floquet topological insulators. *Nature*, 496(7444):196–200, April 2013.

[17] Tim Sanchez, Daniel T. N. Chen, Stephen J. DeCamp, Michael Heymann, and Zvonimir Dogic. Spontaneous motion in hierarchically assembled active matter. *Nature*, 491(7424):431–434, November 2012.

[18] Daniel M. Sussman, Olaf Stenull, and T. C. Lubensky. Topological boundary modes in jammed matter. *Soft Matter*, 12(28):6079–6087, 2016. arXiv: 1512.04480.

[19] Roman Süsstrunk and Sebastian D. Huber. Observation of phononic helical edge states in a mechanical 'topological insulator'. *arXiv preprint*, 1503.06808 [cond-mat], March 2015. arXiv: 1503.06808.

[20] T. Thonhauser and David Vanderbilt. Insulator/Chern-insulator transition in the Haldane model. *Physical Review B*, 74(23):235111, December 2006.

[21] M. F. Thorpe. Continuous deformations in random networks. *J Non Cryst Solids*, 57(3):355–370, September 1983.

[22] D. J. Thouless. Wannier functions for magnetic subbands. *Journal of Physics C: Solid State Physics*, 17(12):L325, 1984.

[23] Pai Wang, Ling Lu, and Katia Bertoldi. Topological Phononic Crystals with One-Way Elastic Edge Waves. *Physical Review Letters*, 115(10), September 2015.

[24] D. Weaire. Existence of a Gap in the Electronic Density of States of a Tetrahedrally Bonded Solid of Arbitrary Structure. *Physical Review Letters*, 26(25):1541–1543, June 1971.

[25] D. Weaire and M. F. Thorpe. Electronic Properties of an Amorphous Solid. I. A Simple Tight-Binding Theory. *Physical Review B*, 4(8):2508–2520, October 1971.